

QUT Digital Repository:  
<http://eprints.qut.edu.au/>



This is the accepted version of this article. Published as:

Frost, Ray L. and Bahfenne, Silmarilly and Keeffe, Eloise C. (2009) *Raman spectroscopic study of the antimonate mineral brandholzite  $Mg[Sb(OH)_6].6H_2O$* . Journal of Raman Spectroscopy, 40(12). pp. 1907-1910.

© Copyright 2009 John Wiley & Sons, Ltd.

The definitive version is available at [www3.interscience.wiley.com](http://www3.interscience.wiley.com)

**Raman spectroscopic study of the antimonate mineral brandholzite  
Mg[Sb(OH)<sub>6</sub>] $\cdot$ 6H<sub>2</sub>O**

**Ray L. Frost,<sup>1</sup> • Jiří Čejka,<sup>1,2</sup> Jiří Sejkora,<sup>2</sup>  
Daniel Ozdín,<sup>3</sup> Silmarilly Bahfenne and Eloise C. Keeffe<sup>1</sup>**

<sup>1</sup> Inorganic Materials Research Program, School of Physical and Chemical Sciences, Queensland University of Technology, GPO Box 2434, Brisbane Queensland 4001, Australia.

<sup>2</sup> National Museum, Václavské náměstí 68, CZ-115 79 Praha 1, Czech Republic.

<sup>3</sup> Department of Mineralogy and Petrology, Faculty of Science, Comenius University, Mlynská Dolina, SK-84 215, Bratislava, Slovak Republic

**Raman spectra of brandholzite Mg[Sb(OH)<sub>6</sub>] $\cdot$ 6H<sub>2</sub>O were studied, complemented with infrared spectra, and related to the structure of the mineral. An intense Raman sharp band at 618 cm<sup>-1</sup> is attributed to the SbO symmetric stretching mode. The low intensity band at 730 cm<sup>-1</sup> is ascribed to the SbO antisymmetric stretching vibration. Low intensity Raman bands were found at 503, 526 and 578 cm<sup>-1</sup>. Corresponding infrared bands were observed at 527, 600, 637, 693, 741 and 788 cm<sup>-1</sup>. Four Raman bands observed at 1043, 1092, 1160 and 1189 cm<sup>-1</sup> and eight infrared bands at 963, 1027, 1055, 1075, 1108, 1128, 1156 and 1196 cm<sup>-1</sup> are assigned to  $\delta$  SbOH deformation modes. A complex pattern resulting from the overlapping band of the water and hydroxyl units is observed. Raman bands are observed at 3240, 3383, 3466, 3483 and 3552 cm<sup>-1</sup>, infrared bands at 3248, 3434 and 3565 cm<sup>-1</sup>. The first two Raman bands and the first infrared band are assigned to water stretching vibrations. The two higher wavenumber Raman bands observed at 3466 and 3552 cm<sup>-1</sup> and two infrared bands at 3434 and 3565 cm<sup>-1</sup> are assigned to the stretching vibrations of the hydroxyl units. Observed Raman and infrared bands are connected with O-H...O hydrogen bonds and their lengths 2.72, 2.79, 2.86, 2.88 and 3.0 Å (Raman) and 2.73, 2.83 and 3.07 Å (infrared).**

**Keywords:** brandholzite, antimonate, antimonite, molecular water, Raman, infrared, spectroscopy

## **Introduction**

The mineral brandholzite, trigonal MgSb<sub>2</sub>(OH)<sub>12</sub> $\cdot$ 6H<sub>2</sub>O, was originally described as plate-like crystals up to 1 mm in size in association with stibnite from mining district Brandholz-Goldkronach (Bavaria, Germany)<sup>1</sup>. Later, it was also

---

• Author to whom correspondence should be addressed (r.frost@qut.edu.au)

found as tiny colorless aggregates at the Krížnica mine, the Pernek deposit, the Malé Karpaty Mountains, western Slovakia, Slovak Republic<sup>2,3</sup> in association with stibnite, sulphur, aragonite, gypsum and antimony-ochres. The structure of the mineral has been published<sup>1,4</sup>. Its crystal structure is isotypic with bottinoite<sup>5,6</sup>. The mineral is isotypic with synthetic  $\text{Co}(\text{H}_2\text{O})_6[\text{Sb}(\text{OH})_6]_2$ <sup>4</sup> and bottinoite,  $\text{Ni}(\text{H}_2\text{O})_6[\text{Sb}(\text{OH})_6]_2$ <sup>5-7</sup>.

Farmer reported the infrared spectra of some synthetic antimonite minerals (see page 413 and 414 with Tables 17. XVIII and XIX)<sup>8</sup>. For the synthetic compound  $\text{NaSb}(\text{OH})_6$  which is a compound with an octahedral structure, infrared bands were observed at 600 and 628  $\text{cm}^{-1}$  (very intense), 735, 775  $\text{cm}^{-1}$  (medium intensity), and 528 and 586  $\text{cm}^{-1}$ . Siebert researched the infrared spectra of selected synthetic antimonates<sup>9,10</sup>. Siebert assigned bands in the 528 to 775  $\text{cm}^{-1}$  region to the stretching vibrations of  $\text{SbO}$  units; in the 1030 to 1120  $\text{cm}^{-1}$  to the deformation modes of  $\text{SbOH}$  units and in the 3220 to 3400  $\text{cm}^{-1}$  to the stretching bands of  $\text{SbOH}$  and water units.

It is interesting to note that only very few papers have been published on the spectroscopy of antimonate minerals. What research has been published is related to the analysis of pigments<sup>11-13</sup>. Some spectroscopic studies of calcium and lead antimonates have been forthcoming<sup>14-16</sup>. Very few studies of related minerals such as mineral antimonates have not been undertaken<sup>17-19</sup>. It was found that the hydroxyl unit was coordinated directly to the metal ion and formed hydrogen bonds to the arsenate anion<sup>20</sup>. Raman spectroscopy has proven especially useful for the study of related minerals<sup>21-29</sup>. As part of a comprehensive study of the molecular structure of secondary minerals containing oxy-anions<sup>30-40</sup>, formed in the oxide zone, using IR and Raman spectroscopy, we report the Raman properties of the antimonate mineral brandholzite. The spectra are related to the mineral structure.

## Experimental

### Mineral

The studied sample of the mineral brandholzite was found at Krížnica mine, the Pernek deposit, the Malé Karpaty Mountains, western Slovakia, Slovak Republic [39], and is deposited in the mineralogical collections of the National Museum Prague. The sample was analysed for phase purity by X-ray powder diffraction. No minor significant impurities were found. Its refined unit-cell parameters for trigonal space group  $P3$ ,  $a$  16.1076(9),  $c$  9.8628(9) Å,  $V$  2216.1(2) Å<sup>3</sup>, are comparable with the data published for this mineral phase<sup>1</sup>. The mineral was analysed by electron microprobe (Cameca SX100, WD mode) for chemical composition. The results (mean of 2 point analyses, recalculation to 100 wt. %) are MgO 6.11, CaO 0.52, FeO 0.16,  $\text{Sb}_2\text{O}_5$  55.91,  $\text{H}_2\text{O}$  37.30 wt. %, sum 100.00 wt. %. The water content was calculated on the basis of theoretical content ( $\text{OH}$ ) = 12 and  $\text{H}_2\text{O}$  = 6 *pfu*. The resulting empirical formula is  $(\text{Mg}_{0.88}\text{Ca}_{0.06}\text{Fe}_{0.01})_{\Sigma 0.95}(\text{H}_2\text{O})_{6.00}[\text{Sb}(\text{OH})_6]_{\Sigma 2.01}$ .

### Raman spectroscopy

The crystals of brandholzite were placed and oriented on the stage of an Olympus BHSM microscope, equipped with 10x and 50x objectives and part of a

Renishaw 1000 Raman microscope system, which also includes a monochromator, a filter system and a Charge Coupled Device (CCD). Further details have been published<sup>21-29</sup>.

## IR spectroscopy

The FTIR spectrum of brandholzite was obtained with the FTIR Nicolet 740 spectrometer using the conventional KBr-disk technique. Infrared spectrum in the range 4000-400  $\text{cm}^{-1}$  was obtained by the co-addition of 32 scans with a resolution of 2  $\text{cm}^{-1}$  and a mirror velocity of 0.1496  $\text{cm/s}$ . Spectral manipulation such as baseline adjustment, smoothing and normalization were performed using the OMNIC software package (Thermo Electron Corporation). Band component analysis was undertaken using the same software package which enabled the type of fitting function to be selected and allows specific parameters to be fixed or varied accordingly. Band fitting was done using a Lorentz-Gauss cross-product function with the minimum number of component bands used for the fitting process which was undertaken until reproducible results were obtained with minimum value of standard errors (usually lower than 2, the range for brandholzite spectra is 1.30-0.20).

## Results and discussion

### Raman spectroscopy

The Raman spectrum of brandholzite in the 400 to 800  $\text{cm}^{-1}$  region is shown in Fig. 1. The spectrum shows some complexity with a number of overlapping bands observed at 618, 630 and 730  $\text{cm}^{-1}$  with additional low intensity bands found at 503, 526 and 578  $\text{cm}^{-1}$ . Infrared bands were observed at 527, 600, 637, 693, 741 and 788  $\text{cm}^{-1}$  (Supplementary Figure 1S). According to Siebert<sup>9,10</sup> all bands in these positions are assignable to stretching vibrations. The observation of multiple bands in the Raman spectrum provides evidence for the non-equivalence of SbO units in the brandholzite structure. In the infrared spectrum of antimony pentoxide an intense band is observed at 740  $\text{cm}^{-1}$  and low intensity bands at ~370, 450 and 680  $\text{cm}^{-1}$ <sup>41</sup>. The infrared spectrum of valentinite ( $\text{Sb}_2\text{O}_3$ )<sub>4</sub> showed bands in similar positions<sup>41</sup>. Farmer reported the band positions of synthetic antimonates of formula  $\text{MSbO}_4$  where M is Cr, Fe, Ga or Rh with a rutile type structure<sup>8</sup>. As such these structures should have four Raman active bands ( $A_{1g} + B_{1g} + B_{2g} + E_g$ ) and four infrared active bands ( $A_{2u} + 3E_u$ ). Infrared bands were observed in the 660 to 735  $\text{cm}^{-1}$ , 520 to 585  $\text{cm}^{-1}$ , 285 to 375  $\text{cm}^{-1}$  and 170 to 190  $\text{cm}^{-1}$ . Although no assignment was given to these bands but one possible interpretation is that the first band is attributed to the antisymmetric stretching mode, the second to the symmetric stretching mode, the third to bending modes and the fourth to a lattice modes. Hence the bands of brandholzite at around 618  $\text{cm}^{-1}$  are attributable the symmetric stretching modes. The observation of several bands in this region suggests that the  $\text{Sb}(\text{OH})_6^{2-}$  are not equivalent. In the infrared spectrum of the compound  $\text{NaSb}(\text{OH})_6$  which has an octahedral structure, a very intense band is observed at 628  $\text{cm}^{-1}$  with bands of lower intensity at 775 and 528  $\text{cm}^{-1}$ <sup>8</sup>.

The Raman spectrum of brandholzite in the 100 to 400  $\text{cm}^{-1}$  region is displayed in Figure 2. Intense Raman bands are observed at 303, 318 and 340  $\text{cm}^{-1}$ . One likely assignment of these bands is to OSbO bending modes. Such an assessment fits well with the assignment of bands for  $\text{MSbO}_4$  structures as reported by Farmer<sup>8</sup>. The observation of multiple bands suggests the non-equivalence of SbO units in the

brandholzite structure. Other low intensity bands are observed at 115, 147, 191, 232 and 252  $\text{cm}^{-1}$ . Low wavenumber bands are observed for  $\text{MSbO}_4$ .<sup>8</sup>

Raman bands in the 900 to 1800  $\text{cm}^{-1}$  region are reported in Fig. 3. Raman bands in this spectral region are of a very low intensity. (The scaling factor is  $\times 7000$  in Fig. 3) Two sets of bands are observed (a) in the 1000 to 1200  $\text{cm}^{-1}$  region and (b) centred upon 1648  $\text{cm}^{-1}$ . Four Raman bands are observed at 1043, 1092, 1160 and 1189  $\text{cm}^{-1}$ . Eight infrared bands at 963, 1027, 1055, 1075, 1108, 1128, 1156 and 1196  $\text{cm}^{-1}$  were observed (Supplementary information Figure 2S). These bands are assigned to  $\delta$  SbOH deformation modes. Siebert reported four infrared bands at 1030, 1075, 1105 and 1120  $\text{cm}^{-1}$  for the synthetic compound  $\text{NaSb}(\text{OH})_6$ . The position and number of these infrared bands for his compound is in good agreement with the position of the Raman bands of brandholzite. The Raman band at 1648  $\text{cm}^{-1}$  is ascribed to the water HOH bending mode. The position of the band for liquid water is  $\sim 1630 \text{ cm}^{-1}$ . The position of the band at 1648  $\text{cm}^{-1}$  and infrared bands at 1632 and 1653  $\text{cm}^{-1}$  (Supplementary information Figure 3S) provides evidence for the water being strongly hydrogen-bonded in the brandholzite structure. Weak infrared bands at 1541, 1558 and 1739  $\text{cm}^{-1}$  may be probably connected with overtones and/or combination bands.

The Raman spectrum of brandholzite in the 2600 to 3800  $\text{cm}^{-1}$  region is shown in Figure 4. A complex pattern resulting from the overlapping bands of the water and OH units is observed. Raman bands are observed at 3240, 3383, 3466, 3483 and 3552  $\text{cm}^{-1}$ , infrared bands at 3248, 3434 and 3565  $\text{cm}^{-1}$ . The Raman bands at 3240 and 3383  $\text{cm}^{-1}$  and the infrared band at 3248  $\text{cm}^{-1}$  are assigned to water stretching vibrations. The higher wavenumber bands observed at 3466 and 3552  $\text{cm}^{-1}$  (Raman) and 3434 and 3565  $\text{cm}^{-1}$  (infrared) (Supplementary information Figure 4S) are assigned to the stretching vibrations of the OH units. There may be inferred a hydrogen-bonding network in the crystal structure of brandholzite with corresponding O-H...O hydrogen bond lengths<sup>42</sup> 2.72, 2.79, 2.86, 2.88 and 3.0 Å [Raman], and 2.73, 2.83 and 3.07 Å (infrared).

## CONCLUSIONS

Raman spectra of brandholzite  $\text{Mg}[\text{Sb}(\text{OH})_6] \cdot 6\text{H}_2\text{O}$  have been studied and related to the structure of the mineral. Raman bands were assigned SbO stretching modes,  $\delta$  SbOH deformation modes, OH stretching modes and water stretching and bending modes. O-H...O hydrogen bond lengths in the crystal structure of brandholzite were inferred from the Raman and infrared spectra.

## Acknowledgments

This work was supported by Ministry of Culture of the Czech Republic (project MK00002327201) to Jiří Sejkora. The authors thank to Miroslava Novotná (VŠCHT, Prague) for her kind support of recording the IR spectrum of brandholzite.

197 The financial and infra-structure support of the Queensland University of Technology,  
198 Inorganic Materials Research Program is gratefully acknowledged. The Australian  
199 Research Council (ARC) is thanked for funding the instrumentation.  
200  
201

## References

- (1) Friedrich A, Wildner M, Tillmanns E, Merz PL *American Mineralogist* **2000**, 85, 593-599.
- (2) Sejkora J, Ozdín D, Vitáloš J, R. Ďud'a *Journal of Geosciences*.
- (3) Sejkora S, Ozdín D, Vitáloš J, Tuček P, Ďud'a R *Lapis* **2004**, 29, 27-36.
- (4) Friedrich A, Mazzi F, Wildner M, Tillmanns E *American Mineralogist* **2003**, 88, 462-463.
- (5) Bonazzi P, Mazzi F *American Mineralogist* **1996**, 81, 1494-1500.
- (6) Bonazzi P, Menchetti S, Caneschi A, Magnanelli S *American Mineralogist* **1992**, 77, 1301-4.
- (7) Clark AM, Rust SA *Mineralogical Magazine* **1993**, 57, 543-4.
- (8) Farmer VC *Mineralogical Society Monograph 4: The Infrared Spectra of Minerals*, 1974.
- (9) Siebert H Z. *anorg. u. allgem. Chem.* **1959**, 301, 161-70.
- (10) Siebert H *Anwendungen der Schwingungsspektroskopie in der Anorganischen Chemie (Anorganische und Allgemeine Chemie in Einzeldarstellungen, Bd. 7) (Application of Vibrational Spectroscopy in Inorganic Chemistry (Monographs in Inorganic and General Chemistry, Vol. 7))*, 1966.
- (11) Correia AM, Oliveira MJV, Clark RJH, Ribeiro MI, Duarte ML *Analytical Chemistry (Washington, DC, United States)* **2008**, 80, 1482-1492.
- (12) Correia AM, Clark RJH, Ribeiro MIM, Duarte MLTS *Journal of Raman Spectroscopy* **2007**, 38, 1390-1405.
- (13) Clark RJH, Cridland L, Kariuki BM, Harris KDM, Withnall R *Journal of the Chemical Society, Dalton Transactions: Inorganic Chemistry* **1995**, 2577-82.
- (14) Husson E, Repelin Y, Vandenborre MT *Spectrochimica Acta, Part A: Molecular and Biomolecular Spectroscopy* **1984**, 40A, 1017-20.
- (15) Haeuseler H *Spectrochimica Acta, Part A: Molecular and Biomolecular Spectroscopy* **1981**, 37A, 487-95.
- (16) Vandenborre MT, Husson E, Brusset H, Cerez A *Spectrochimica Acta, Part A: Molecular and Biomolecular Spectroscopy* **1980**, 36A, 1045-52.
- (17) Paques-Ledent MT, Tarte P *Spectrochim. Acta, Part A* **1974**, 30A, 673-89.
- (18) Gevork'yan SV, Povarennykh AS *Konst. Svoistva Miner.* **1975**, 9, 73-81.
- (19) Braithwaite RSW *Mineral. Mag.* **1983**, 47, 51-7.
- (20) Sumin De Portilla VI *Can. Mineral.* **1974**, 12, 262-8.
- (21) Frost RL, Cejka J, Ayoko G *Journal of Raman Spectroscopy* **2008**, 39, 495-502.
- (22) Frost RL, Cejka J, Ayoko GA, Dickfos MJ *Journal of Raman Spectroscopy* **2008**, 39, 374-379.
- (23) Frost RL, Cejka J, Dickfos MJ *Journal of Raman Spectroscopy* **2008**, 39, 779-785.
- (24) Frost RL, Dickfos MJ, Cejka J *Journal of Raman Spectroscopy* **2008**, 39, 582-586.
- (25) Frost RL, Hales MC, Wain DL *Journal of Raman Spectroscopy* **2008**, 39, 108-114.
- (26) Frost RL, Keeffe EC *Journal of Raman Spectroscopy* **2008**, in press.
- (27) Frost RL, Locke A, Martens WN *Journal of Raman Spectroscopy* **2008**, 39, 901-908.
- (28) Frost RL, Reddy BJ, Dickfos MJ *Journal of Raman Spectroscopy* **2008**, 39, 909-913.

- 252 (29) Palmer SJ, Frost RL, Ayoko G, Nguyen T *Journal of Raman Spectroscopy*  
 253 **2008**, 39, 395-401.
- 254 (30) Frost RL, Bouzaid JM *Journal of Raman Spectroscopy* **2007**, 38, 873-879.
- 255 (31) Frost RL, Bouzaid JM, Martens WN, Reddy BJ *Journal of Raman*  
 256 *Spectroscopy* **2007**, 38, 135-141.
- 257 (32) Frost RL, Cejka J *Journal of Raman Spectroscopy* **2007**, 38, 1488-1493.
- 258 (33) Frost RL, Cejka J, Ayoko GA, Weier ML *Journal of Raman Spectroscopy*  
 259 **2007**, 38, 1311-1319.
- 260 (34) Frost RL, Cejka J, Weier ML *Journal of Raman Spectroscopy* **2007**, 38, 460-  
 261 466.
- 262 (35) Frost RL, Cejka J, Weier ML, Martens WN, Ayoko GA *Journal of Raman*  
 263 *Spectroscopy* **2007**, 38, 398-409.
- 264 (36) Frost RL, Dickfos MJ *Journal of Raman Spectroscopy* **2007**, 38, 1516-1522.
- 265 (37) Frost RL, Palmer SJ, Bouzaid JM, Reddy BJ *Journal of Raman Spectroscopy*  
 266 **2007**, 38, 68-77.
- 267 (38) Frost RL, Pinto C *Journal of Raman Spectroscopy* **2007**, 38, 841-845.
- 268 (39) Frost RL, Weier ML, Williams PA, Leverett P, Kloprogge JT *Journal of*  
 269 *Raman Spectroscopy* **2007**, 38, 574-583.
- 270 (40) Locke AJ, Martens WN, Frost RL *Journal of Raman Spectroscopy* **2007**, 38,  
 271 1429-1435.
- 272 (41) Gadsden JA *Infrared spectra of minerals and related inorganic compounds*;  
 273 Butterworth: London, UK, 1975.
- 274 (42) Libowitzky E *Monatshefte fuer Chemie* **1999**, 130, 1047-1059.
- 275
- 276
- 277



278 ***List of Figures***

279

280 Fig. 1 Raman spectrum of brandholzite in the 600 to 1200  $\text{cm}^{-1}$  region

281

282 Fig. 2 Raman spectrum of brandholzite in the 100 to 600  $\text{cm}^{-1}$  region

283

284 Fig. 3 Raman spectrum of brandholzite in the 1200 to 1800  $\text{cm}^{-1}$  region

285

286 Fig. 4 Raman spectrum of brandholzite in the 3000 to 3700  $\text{cm}^{-1}$  region

287

288

289 Fig. S1 Infrared spectrum of brandholzite in the 490 to 900  $\text{cm}^{-1}$  region

290


291 Fig. S1 Infrared spectrum of brandholzite in the 910 to 1240  $\text{cm}^{-1}$  region

292


293 Fig. S1 Infrared spectrum of brandholzite in the 1490 to 1800  $\text{cm}^{-1}$  region

294


295 Fig. S1 Infrared spectrum of brandholzite in the 2600 to 3850  $\text{cm}^{-1}$  region




**Figure 1**



**Figure 2**



**Figure 3**



**Figure 4**